

Wavelet Linear Density Estimation for a GARCH Model under Various Dependence Structures

Christophe Chesneau¹, Hassan Doosti²

¹Laboratoire de Mathématiques Nicolas Oresme, Université de Caen Basse-Normandie, Campus II, Science 3, France.

²Department of Mathematics, Tarbiat Moallem University, Tehran, Iran.

Abstract. We consider n observations from the GARCH-type model: $S = \sigma^2 Z$, where σ^2 and Z are independent random variables. We develop a new wavelet linear estimator of the unknown density of σ^2 under four different dependence structures: the strong mixing case, the β -mixing case, the pairwise positive quadrant case and the ρ -mixing case. Its asymptotic mean integrated squared error properties are explored. In each case, we prove that it attains a fast rate of convergence.

Keywords. Dependent sequence, GARCH models, linear estimator, rate of convergence, wavelets.

MSC: 62G07, 62G20.

1 Introduction

The following GARCH-type model is considered: let $(S_i)_{i \in \mathbb{Z}}$ be a strictly stationary random sequence such that

$$S_i = \sigma_i^2 Z_i, \quad i \in \mathbb{Z}, \quad (1)$$

$(Z_i)_{i \in \mathbb{Z}}$ is a sequence of identically distributed random variables with common known density $f_Z : [0, 1] \rightarrow (0, \infty)$ and $(\sigma_i^2)_{i \in \mathbb{Z}}$ is a sequence of

Christophe Chesneau(✉)(chesneau@math.unicaen.fr),

Hassan Doosti (h.doosti@tmu.ac.ir)

Received: February 18, 2011; Accepted: October 30, 2011

identically distributed random variables with common unknown density $f_{\sigma^2} : [0, 1] \rightarrow (0, \infty)$. For any $i \in \mathbb{Z}$, Z_i and σ_i^2 are independent. We aim to estimate f_{σ^2} when only n random variables S_1, \dots, S_n are observed under various dependent structures. Financial applications related to (1) can be found in [5].

In most of the papers, (1) is re-written via a logarithmic transformation: $\ln S_i = \ln \sigma_i^2 + \ln Z_i$, $i \in \mathbb{Z}$. Since we have a sum of two independent random variables, the density of $\ln \sigma_1^2$ becomes a convolution product. The classical scheme consists in deconvolving and estimating this density by using Fourier transform and nonparametric methods. For dependent sequences, see e.g. [21], [10] and [31]. However, the estimation of f_{σ^2} is not “direct” in the following sense: if we denote $f_{\ln \sigma^2}$ the density of $\ln \sigma_1^2$, we have $f_{\sigma^2}(x) = (1/x)f_{\ln \sigma^2}(\ln x)$, $x \in (0, 1)$ and, for any estimator \hat{f} of $f_{\ln \sigma^2}$, the associated mean integrated squared error (MISE) is

$$\mathbb{E} \left(\int_{-\infty}^0 \left(\hat{f}(x) - f_{\ln \sigma^2}(x) \right)^2 dx \right) = \mathbb{E} \left(\int_0^1 \left(\frac{1}{x} \hat{f}(\ln x) - f_{\sigma^2}(x) \right)^2 x dx \right).$$

Thus we obtain the MISE for $(1/x)\hat{f}(\ln x)$ of f_{σ^2} but with respect to the measure $x dx$, not dx . For this reason, the global estimation of f_{σ^2} on $[0, 1]$ from $\ln S_1, \dots, \ln S_n$ via the standard MISE (with respect to dx) is not obvious. This point is also underlined in [10, 3.5]. Considering the exponential strong mixing case, the “direct” estimation of f_{σ^2} has been recently investigated by [8].

In this paper, we complete this last study by estimating “directly” f_{σ^2} for other realistic and standard dependence conditions as the polynomial strong mixing dependence (introduced by [26]), the β -mixing dependence (introduced by [33]), the pairwise positive quadrant dependence (PPQD) (introduced by [20]) and the ρ -mixing dependence (introduced by [18]). For results, examples and references on the standard density estimation problem under such dependence conditions, see e.g. [19], [4], [32], [30], [11], [21], [25], [6] and [17].

Combining the approaches of [8] and [17], we construct an estimator based on wavelet projections. We use wavelets because of their computational efficiency and asymptotic optimality properties. In particular, wavelet estimators enjoy interesting MISE for functions having possible complex singularities. We refer to [2] and [16] for a detailed coverage of wavelet theory in statistics. The asymptotic performance of our estimator is evaluated by determining an upper bound of the MISE over Besov balls. It is obtained as sharp as possible and coincides with the

one related to the standard *i.i.d.* framework.

The organization of the paper is as follows. Assumptions on the model are presented in Section 2. Section 3 describes the wavelet basis and the Besov balls. Section 4 is devoted to our linear wavelet estimator and a general result. Applications are set in Section 5. Technical proofs are collected in Section 6.

2 Assumptions

Set $\mathbb{L}^2([0, 1]) = \left\{ h : [0, 1] \rightarrow \mathbb{R}; \int_0^1 (h(x))^2 dx < \infty \right\}$. We assume that $f_{\sigma^2} \in \mathbb{L}^2([0, 1])$.

We suppose that there exists a positive integer ν such that, for any $i \in \{1, \dots, n\}$,

$$Z_i = \prod_{r=1}^{\nu} U_{r,i}, \quad (2)$$

where $U_{1,i}, \dots, U_{\nu,i}$ are ν *i.i.d.* random variables having the common uniform distribution on $[0, 1]$. Assumption (2) excludes the Gaussian case but occurs in the study of some GARCH-type model as, for instance, the generalized multiplicative censoring model (see e.g. [1]).

It follows from (2) that

- the density of Z_1 is

$$f_Z(x) = \frac{1}{(\nu - 1)!} (-\ln x)^{\nu-1}, \quad x \in [0, 1].$$

- the density of S_1 is

$$\begin{aligned} f_S(x) &= \int_x^1 f_Z\left(\frac{x}{y}\right) f_{\sigma^2}(y) \frac{1}{y} dy \\ &= \frac{1}{(\nu - 1)!} \sum_{u=0}^{\nu-1} \binom{\nu-1}{u} (-\ln x)^u \int_x^1 (\ln y)^{\nu-1-u} f_{\sigma^2}(y) \frac{1}{y} dy, \\ &\quad x \in [0, 1]. \end{aligned} \quad (3)$$

3 Wavelets and Besov Balls

For the purposes of this paper, we use the compactly supported wavelet bases on $[0, 1]$ described below.

Let N be an integer such that $N > \nu + 1$ (where ν is the one in (2)), ϕ and ψ be the initial wavelets of the Daubechies wavelets $db2N$. These functions have the particularity to be compactly supported and to belong to the class \mathcal{C}^ν .

Set

$$\phi_{j,k}(x) = 2^{j/2}\phi(2^j x - k), \quad \psi_{j,k}(x) = 2^{j/2}\psi(2^j x - k).$$

With an appropriate treatment at the boundaries, there exists an integer τ satisfying $2^\tau \geq 2N$ such that the collection

$$\mathcal{B} = \{\phi_{\tau,k}(\cdot), k \in \{0, \dots, 2^\tau - 1\}; \psi_{j,k}(\cdot); \\ j \in \mathbb{N} - \{0, \dots, \tau - 1\}, k \in \{0, \dots, 2^j - 1\}\},$$

is an orthonormal basis of $\mathbb{L}^2([0, 1])$. We refer to [9].

For any integer $\ell \geq \tau$, any $h \in \mathbb{L}^2([0, 1])$ can be expanded on \mathcal{B} as

$$h(x) = \sum_{k=0}^{2^\ell-1} \alpha_{\ell,k} \phi_{\ell,k}(x) + \sum_{j=\ell}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k} \psi_{j,k}(x), \quad x \in [0, 1],$$

where $\alpha_{j,k}$ and $\beta_{j,k}$ are the wavelet coefficients of h defined by

$$\alpha_{j,k} = \int_0^1 h(x) \phi_{j,k}(x) dx, \quad \beta_{j,k} = \int_0^1 h(x) \psi_{j,k}(x) dx. \quad (4)$$

As is traditional in the wavelet estimation literature, we shall investigate the performances of our estimator by assuming that f_{σ^2} belongs to Besov balls. Their definitions in terms of wavelet coefficients are given below.

Let $s > 0$ and $M > 0$. A function h belongs to $B_{2,\infty}^s(M)$ if and only if there exists a constant $M^* > 0$ (depending on M) such that the associated wavelet coefficients (4) satisfy

$$\sup_{j \geq \tau} 2^{2js} \sum_{k=0}^{2^j-1} \beta_{j,k}^2 \leq M^*.$$

The Besov balls capture a wide variety of smoothness features in a function. Further details can be found in [22].

4 Estimator and result

4.1 Linear wavelet estimator

We follow the methodology of [8, Lemma 1]. For any positive integer ℓ and any $h \in \mathcal{C}^\ell([0, 1])$, set

$$T(h)(x) = (xh(x))', \quad T_\ell(h)(x) = T(T_{\ell-1}(h))(x), \quad x \in [0, 1]. \quad (5)$$

The definition of this operator is such that, for any $h \in \mathcal{C}^\nu([0, 1])$, we have

$$\int_0^1 f_{\sigma^2}(x)h(x)dx = \int_0^1 f_S(x)T_\nu(h)(x)dx. \quad (6)$$

The proof of (6) is based on the two following steps:

Step 1. For any positive integer ℓ , set $G(h)(x) = -xh'(x)$ and $G_\ell(h)(x) = G(G_{\ell-1}(h))(x)$. It follows from (3), derivations and the binomial theorem that

$$f_{\sigma^2}(x) = -x(G_{\nu-1}(f_S)(x))' = G_\nu(f_S)(x), \quad x \in [0, 1]. \quad (7)$$

Step 2. By (7) and ν integrations by parts, we have

$$\begin{aligned} \int_0^1 f_{\sigma^2}(x)h(x)dx &= \int_0^1 G_\nu(f_S)(x)h(x)dx \\ &= \int_0^1 G_{\nu-1}(f_S)(x)T(h)(x)dx \\ &= \dots = \int_0^1 f_S(x)T_\nu(h)(x)dx. \end{aligned}$$

Note that, in the simplest case $\nu = 1$, since $f_{\sigma^2}(x) = -xf'_S(x)$, $x \in [0, 1]$, the proof is reduced to

$$\begin{aligned} \int_0^1 f_{\sigma^2}(x)h(x)dx &= -\int_0^1 f'_S(x)xh(x)dx = \int_0^1 f_S(x)(xh(x))'dx \\ &= \int_0^1 f_S(x)T(h)(x)dx. \end{aligned}$$

Using the method of moments, for any integer $j \geq \tau$ and any $k \in \{0, \dots, 2^j - 1\}$, we estimate the unknown wavelet coefficient $\alpha_{j,k} = \int_0^1 f_{\sigma^2}(x)\phi_{j,k}(x)dx$ by

$$\hat{\alpha}_{j,k} = \frac{1}{n} \sum_{i=1}^n T_\nu(\phi_{j,k})(S_i),$$

(ν is the one in (2)).

Assuming that $f_{\sigma^2} \in B_{2,\infty}^s(M)$, we define the linear estimator \hat{f} by

$$\hat{f}(x) = \sum_{k=0}^{2^{j_0}-1} \hat{\alpha}_{j_0,k} \phi_{j_0,k}(x), \quad x \in [0, 1], \quad (8)$$

where j_0 is an integer which will be chosen later (see Theorem 4.1 below).

Such an estimator is standard in nonparametric estimation via wavelets. See e.g. [16, Section 10]. For a survey on wavelet linear estimators in various density models, we refer to [7].

4.2 Result

Theorem 4.1. *Consider (1) under the assumptions of Section 2. We suppose that*

- *there exist three constants, $\gamma \geq 2\nu$, $\delta \in [0, 1)$ and $C > 0$, such that, for any integer $j \geq \tau$,*

$$\sum_{k=0}^{2^j-1} \sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_m), T_\nu(\phi_{j,k})(S_0))| \leq C 2^{j(\gamma+1)} n^\delta, \quad (9)$$

where $\mathbb{C}_{ov}(\cdot, \cdot)$ denotes the covariance function and T_ν is (5).

- *there exists a constant $C_* > 0$ such that*

$$\sup_{x \in [0, 1]} f_S(x) \leq C_*, \quad (10)$$

Suppose that $f_{\sigma^2} \in B_{2,\infty}^s(M)$ with $s > 0$ and $M > 0$. Let \hat{f} be (8) with j_0 such that $2^{j_0} = \lceil n^{(1-\delta)/(2s+\gamma+1)} \rceil$ (where $\lceil a \rceil$ denotes the integer part of a). Then there exists a constant $C > 0$ such that

$$\mathbb{E} \left(\int_0^1 (\hat{f}(x) - f_{\sigma^2}(x))^2 dx \right) \leq C n^{-2s(1-\delta)/(2s+\gamma+1)}.$$

Naturally, the rate of convergence in Theorem 4.1 is obtained to be as sharp as possible.

The assumption (9) measures the dependence between S_m and S_0 . It is close to the Ξ -weak dependence introduced by [13, 14] but with the operator T_ν .

5 Applications

The four following subsections investigate separately the strong mixing case, the β -mixing case, the PPQD case and the ρ -dependence case which occur in a large variety of applications. We refer to [35], [27], [24] and [3].

5.1 Application to the strong mixing dependence

Definition 5.1. Let $(Y_i)_{i \in \mathbb{Z}}$ be a strictly stationary random sequence. For any $m \in \mathbb{Z}$, we define the m -th strong mixing coefficient of $(Y_i)_{i \in \mathbb{Z}}$ by

$$\alpha_m = \sup_{(A,B) \in \mathcal{F}_{-\infty,0}^Y \times \mathcal{F}_{m,\infty}^Y} |\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)|,$$

where $\mathcal{F}_{-\infty,0}^Y$ is the σ -algebra generated by \dots, Y_{-1}, Y_0 and $\mathcal{F}_{m,\infty}^Y$ is the σ -algebra generated by Y_m, Y_{m+1}, \dots .

We say that $(Y_i)_{i \in \mathbb{Z}}$ is strong mixing if and only if $\lim_{m \rightarrow \infty} \alpha_m = 0$.

Applications on strong mixing can be found in e.g. [34], [12] and [5]. In the context of GARCH-type models as (1), see [15].

Proposition 5.1. Consider (1) under the assumptions of Section 2. Suppose that

- $(S_i)_{i \in \mathbb{Z}}$ is strong mixing,
- there exist three constants, $q \in (0, 1)$, $\delta \in [0, 1)$ and $C > 0$, such that

$$\sum_{m=1}^n m^q \alpha_m^q \leq Cn^\delta, \quad (11)$$

- there exists a constant $C_* > 0$ such that

$$\sup_{m \in \{1, \dots, n\}} \sup_{(x,y) \in [0,1]^2} f_{(S_m, S_0)}(x, y) \leq C_*, \quad \sup_{x \in [0,1]} f_S(x) \leq C_*, \quad (12)$$

where $f_{(S_m, S_0)}$ is the density of (S_m, S_0) and f_S is (3).

Suppose that $f_{\sigma^2} \in B_{2,\infty}^s(M)$ with $s > 0$ and $M > 0$. Let \hat{f} be (8) with j_0 such that $2^{j_0} = \lceil n^{(1-\delta)/(2s+2\nu+1)} \rceil$. Then there exists a constant $C > 0$ such that

$$\mathbb{E} \left(\int_0^1 \left(\hat{f}(x) - f_{\sigma^2}(x) \right)^2 dx \right) \leq Cn^{-2s(1-\delta)/(2s+2\nu+1)}.$$

Note that (11) includes strong mixing coefficients with a polynomial rate of decay. For instance, if $\alpha_m = 1/m^p$ with $p > 1$, then (11) holds with $\delta = 0$ (by taking $q \in (1/(p-1), 1)$).

The first inequality in (12) can be viewed as a ‘‘Castellana-Leadbetter’’-type condition. It is standard in nonparametric estimation via dependent observations. Remark that, thanks to this assumption, Proposition 5.1 improves [8, Theorem1]: the obtained rate of convergence is faster; the parameter q does not deteriorate the rate of convergence. However, this condition seems difficult to check in some situations. An alternative is explored in the next subsection.

5.2 Application to the β -mixing dependence

Definition 5.2. Let $(Y_i)_{i \in \mathbb{Z}}$ be a strictly stationary random sequence. For any $m \in \mathbb{Z}$, we define the m -th β -mixing coefficient of $(Y_i)_{i \in \mathbb{Z}}$ by

$$\beta_m = \frac{1}{2} \sup_{((A_i)_{i \in I}, (B_j)_{j \in J}) \in \mathcal{F}_{-\infty, 0}^Y \times \mathcal{F}_{m, \infty}^Y} \sum_{i \in I} \sum_{j \in J} |\mathbb{P}(A_i \cap B_j) - \mathbb{P}(A_i)\mathbb{P}(B_j)|,$$

where the supremum is taken over all finite partitions $(A_i)_{i \in I}$ and $(B_j)_{j \in J}$ of Ω , which are respectively $\mathcal{F}_{-\infty, 0}^Y$ and $\mathcal{F}_{m, \infty}^Y$ measurable, $\mathcal{F}_{-\infty, 0}^Y$ is the σ -algebra generated by \dots, Y_{-1}, Y_0 and $\mathcal{F}_{m, \infty}^Y$ is the one generated by Y_m, Y_{m+1}, \dots .

We say that $(Y_i)_{i \in \mathbb{Z}}$ is β -mixing if and only if $\lim_{m \rightarrow \infty} \beta_m = 0$.

Full details can be found in e.g. [12], [32], [5] and [3].

Proposition 5.2. Consider (1) under the assumptions of Section 2. Suppose that

- $(S_i)_{i \in \mathbb{Z}}$ is β -mixing,
- there exists a constant $C > 0$ such that

$$\sum_{m=1}^{\infty} \beta_m \leq C, \tag{13}$$

- there exists a constant $C_* > 0$ such that

$$\sup_{x \in [0, 1]} f_S(x) \leq C_*.$$

Suppose that $f_{\sigma^2} \in B_{2,\infty}^s(M)$ with $s > 0$ and $M > 0$. Let \hat{f} be (8) with j_0 such that $2^{j_0} = \lceil n^{1/(2s+2\nu+1)} \rceil$. Then there exists a constant $C > 0$ such that

$$\mathbb{E} \left(\int_0^1 \left(\hat{f}(x) - f_{\sigma^2}(x) \right)^2 dx \right) \leq C n^{-2s/(2s+2\nu+1)}.$$

Since β -mixing implies strong mixing, Proposition 5.2 shows that the rate of convergence $n^{-2s/(2s+2\nu+1)}$ can be attained by \hat{f} for strong mixing $(S_i)_{i \in \mathbb{Z}}$ without the constraint on $f_{(S_m, S_0)}$ in (12).

5.3 Application to the PPQD

Definition 5.3. We say that n random variables S_1, \dots, S_n are pairwise positive quadrant dependent (PPQD) if and only if, for any $(\ell, v) \in \{1, \dots, n\}^2$ with $\ell \neq v$ and any $(x, y) \in [0, 1]^2$,

$$\mathbb{P}(S_\ell > x, S_v > y) \geq \mathbb{P}(S_\ell > x)\mathbb{P}(S_v > y).$$

Further details on PPQD can be found in [20], [23] and [28].

Proposition 5.3. Consider (1) under the assumptions of Section 2. Suppose that

- S_1, \dots, S_n are PPQD,
- there exist two constants, $\delta \in [0, 1)$ and $C > 0$, such that

$$\sum_{m=1}^n m^3 \mathbb{C}_{ov}(S_m, S_0) \leq C n^\delta, \quad (14)$$

- (12) is satisfied.

Suppose that $f_{\sigma^2} \in B_{2,\infty}^s(M)$ with $s > 0$ and $M > 0$. Let \hat{f} be (8) with j_0 such that $2^{j_0} = \lceil n^{(1-\delta)/(2s+2\nu+1)} \rceil$. Then there exists a constant $C > 0$ such that

$$\mathbb{E} \left(\int_0^1 \left(\hat{f}(x) - f_{\sigma^2}(x) \right)^2 dx \right) \leq C n^{-2s(1-\delta)/(2s+2\nu+1)}.$$

Proposition 5.4 below investigates the MISE properties of \hat{f} in the PPQD case without the constraint on $f_{(S_m, S_0)}$ in (12).

Proposition 5.4. Consider (1) under the assumptions of Section 2. Suppose that

- S_1, \dots, S_n are PPQD,
- there exist two constants, $\delta \in [0, 1)$ and $C > 0$, such that

$$\sum_{m=1}^n \mathbb{C}_{ov}(S_m, S_0) \leq Cn^\delta, \quad (15)$$

- there exists a constant $C_* > 0$ such that

$$\sup_{x \in [0, 1]} f_S(x) \leq C_*.$$

Suppose that $f_{\sigma^2} \in B_{2, \infty}^s(M)$ with $s > 0$ and $M > 0$. Let \hat{f} be (8) with j_0 such that $2^{j_0} = \lceil n^{(1-\delta)/(2s+2\nu+4)} \rceil$. Then there exists a constant $C > 0$ such that

$$\mathbb{E} \left(\int_0^1 \left(\hat{f}(x) - f_{\sigma^2}(x) \right)^2 dx \right) \leq Cn^{-2s(1-\delta)/(2s+2\nu+4)}.$$

5.4 Application to the ρ -mixing dependence

Definition 5.4. Let $(Y_i)_{i \in \mathbb{Z}}$ be a strictly stationary random sequence. For any $m \in \mathbb{Z}$, we define the m -th maximal correlation coefficient of $(Y_i)_{i \in \mathbb{Z}}$ by

$$\rho_m = \sup_{(U, V) \in \mathbb{L}^2(\mathcal{F}_{-\infty, 0}^Y) \times \mathbb{L}^2(\mathcal{F}_{m, \infty}^Y)} \frac{|\mathbb{C}_{ov}(U, V)|}{\sqrt{\mathbb{V}(U)\mathbb{V}(V)}},$$

where $\mathcal{F}_{-\infty, 0}^Y$ is the σ -algebra generated by \dots, Y_{-1}, Y_0 , $\mathcal{F}_{m, \infty}^Y$ is the one generated by Y_m, Y_{m+1}, \dots and, for any $\mathcal{A} \in \{\mathcal{F}_{-\infty, 0}^Y, \mathcal{F}_{m, \infty}^Y\}$, $\mathbb{L}^2(\mathcal{A}) = \{U \in \mathcal{A}; \mathbb{E}(U^2) < \infty\}$.

We say that $(Y_i)_{i \in \mathbb{Z}}$ is ρ -mixing if and only if $\lim_{m \rightarrow \infty} \rho_m = 0$.

For details on ρ -mixing, we refer to [18], [12], [29], [19] and [35].

Proposition 5.5. Consider (1) under the assumptions of Section 2. Suppose that

- $(S_i)_{i \in \mathbb{Z}}$ is ρ -mixing,
- there exist two constants, $\delta \in [0, 1)$ and $C > 0$, such that

$$\sum_{m=1}^n \rho_m \leq Cn^\delta, \quad (16)$$

- there exists a constant $C_* > 0$ such that

$$\sup_{x \in [0,1]} f_S(x) \leq C_*. \quad (17)$$

Suppose that $f_{\sigma^2} \in B_{2,\infty}^s(M)$ with $s > 0$ and $M > 0$. Let \hat{f} be (8) with j_0 such that $2^{j_0} = \lceil n^{(1-\delta)/(2s+2\nu+1)} \rceil$. Then there exists a constant $C > 0$ such that

$$\mathbb{E} \left(\int_0^1 \left(\hat{f}(x) - f_{\sigma^2}(x) \right)^2 dx \right) \leq C n^{-2s(1-\delta)/(2s+2\nu+1)}.$$

General Remark. Note that, in Propositions 5.1, 5.3 and 5.5, if $\delta = 0$, the rate of convergence $n^{-2s/(2s+2\nu+1)}$ becomes the one attained by \hat{f} when S_1, \dots, S_n are *i.i.d.*. Therefore, our results extend the good asymptotic performances of \hat{f} in the standard *i.i.d.* case to the considered dependence structures.

Conclusions and Perspectives. We have constructed a new wavelet estimator to estimate a density in a GARCH-type model under various dependence structures. Its asymptotic MISE properties have been investigated and fast rates of convergence have been established.

Due to its construction, \hat{f} is not adaptive with respect to s , ν and δ . Adaptivity can perhaps be achieved by using another wavelet estimator as the hard thresholding one. However, several important technical difficulties arise due to the dependence conditions and it is not immediately clear how to solve them. This aspect needs further investigations that we leave for a future work.

6 Proofs

In this section, we consider (1) under the assumptions of Section 2. Moreover, C denotes any constant that does not depend on j , k and n . Its value may change from one term to another and may depend on ϕ .

Proof of Theorem 4.1. We expand the function f_{σ^2} on \mathcal{B} as

$$f_{\sigma^2}(x) = \sum_{k=0}^{2^{j_0}-1} \alpha_{j_0,k} \phi_{j_0,k}(x) + \sum_{j=j_0}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k} \psi_{j,k}(x), \quad x \in [0, 1],$$

where $\alpha_{j_0,k} = \int_0^1 f_{\sigma^2}(x) \phi_{j_0,k}(x) dx$ and $\beta_{j,k} = \int_0^1 f_{\sigma^2}(x) \psi_{j,k}(x) dx$.

We have, for any $x \in [0, 1]$,

$$\widehat{f}(x) - f_{\sigma^2}(x) = \sum_{k=0}^{2^{j_0}-1} (\widehat{\alpha}_{j_0,k} - \alpha_{j_0,k}) \phi_{j_0,k}(x) - \sum_{j=j_0}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k} \psi_{j,k}(x).$$

Since \mathcal{B} is an orthonormal basis of $\mathbb{L}^2([0, 1])$, we have

$$\begin{aligned} \mathbb{E} \left(\int_0^1 (\widehat{f}(x) - f_{\sigma^2}(x))^2 dx \right) = \\ \sum_{k=0}^{2^{j_0}-1} \mathbb{E} \left((\widehat{\alpha}_{j_0,k} - \alpha_{j_0,k})^2 \right) + \sum_{j=j_0}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k}^2. \end{aligned} \quad (18)$$

Let us now bound these two terms in turn.

Using (6), we have

$$\begin{aligned} \alpha_{j_0,k} &= \int_0^1 f_{\sigma^2}(x) \phi_{j_0,k}(x) dx = \int_0^1 f_S(x) T_{\nu}(\phi_{j_0,k})(x) dx \\ &= \mathbb{E}(T_{\nu}(\phi_{j_0,k})(S_0)) = \mathbb{E}(\widehat{\alpha}_{j_0,k}). \end{aligned}$$

So

$$\mathbb{E} \left((\widehat{\alpha}_{j_0,k} - \alpha_{j_0,k})^2 \right) = \mathbb{V}(\widehat{\alpha}_{j_0,k}) = \frac{1}{n^2} \mathbb{V} \left(\sum_{i=1}^n T_{\nu}(\phi_{j_0,k})(S_i) \right). \quad (19)$$

We have

$$\begin{aligned} &\mathbb{V} \left(\sum_{i=1}^n T_{\nu}(\phi_{j_0,k})(S_i) \right) \\ &= \sum_{v=1}^n \sum_{\ell=1}^n \mathbb{C}_{ov} (T_{\nu}(\phi_{j_0,k})(S_v), T_{\nu}(\phi_{j_0,k})(S_{\ell})) \\ &= n \mathbb{V}(T_{\nu}(\phi_{j_0,k})(S_0)) + 2 \sum_{v=2}^n \sum_{\ell=1}^{v-1} \mathbb{C}_{ov} (T_{\nu}(\phi_{j_0,k})(S_v), T_{\nu}(\phi_{j_0,k})(S_{\ell})) \\ &\leq n \mathbb{V}(T_{\nu}(\phi_{j_0,k})(S_0)) + 2 \left| \sum_{v=2}^n \sum_{\ell=1}^{v-1} \mathbb{C}_{ov} (T_{\nu}(\phi_{j_0,k})(S_v), T_{\nu}(\phi_{j_0,k})(S_{\ell})) \right|. \end{aligned} \quad (20)$$

The stationarity of $(S_i)_{i \in \mathbb{Z}}$ implies that

$$\begin{aligned}
& \left| \sum_{v=2}^n \sum_{\ell=1}^{v-1} \mathbb{C}_{ov} (T_\nu(\phi_{j_0,k})(S_v), T_\nu(\phi_{j_0,k})(S_\ell)) \right| \\
&= \left| \sum_{m=1}^n (n-m) \mathbb{C}_{ov} (T_\nu(\phi_{j_0,k})(S_m), T_\nu(\phi_{j_0,k})(S_0)) \right| \\
&\leq n \sum_{m=1}^n |\mathbb{C}_{ov} (T_\nu(\phi_{j_0,k})(S_m), T_\nu(\phi_{j_0,k})(S_0))|. \tag{21}
\end{aligned}$$

It follows from (19), (20) and (21) that

$$\begin{aligned}
& \mathbb{E} \left((\hat{\alpha}_{j_0,k} - \alpha_{j_0,k})^2 \right) \\
&\leq C \frac{1}{n} \left(\mathbb{V} (T_\nu(\phi_{j_0,k})(S_0)) + \sum_{m=1}^n |\mathbb{C}_{ov} (T_\nu(\phi_{j_0,k})(S_m), T_\nu(\phi_{j_0,k})(S_0))| \right).
\end{aligned}$$

Using [8, Proposition 1] i.e. thanks to (10), $\mathbb{V} (T_\nu(\phi_{j_0,k})(S_0)) \leq C2^{2\nu j_0}$, (9) and $\gamma \geq 2\nu$, we have

$$\begin{aligned}
\sum_{k=0}^{2^{j_0}-1} \mathbb{E} \left((\hat{\alpha}_{j_0,k} - \alpha_{j_0,k})^2 \right) &\leq C \frac{1}{n} \left(2^{j_0} 2^{2\nu j_0} + 2^{j_0(\gamma+1)} n^\delta \right) \\
&\leq C 2^{(\gamma+1)j_0} n^{\delta-1} \\
&\leq C n^{-2s(1-\delta)/(2s+\gamma+1)}. \tag{22}
\end{aligned}$$

Using $f_{\sigma^2} \in B_{2,\infty}^s(M)$, we obtain

$$\sum_{j=j_0}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k}^2 \leq C \sum_{j=j_0}^{\infty} 2^{-2js} \leq C 2^{-2j_0s} \leq C n^{-2s(1-\delta)/(2s+\gamma+1)}. \tag{23}$$

It follows from (18), (22) and (23) that

$$\mathbb{E} \left(\int_0^1 (\hat{f}(x) - f_{\sigma^2}(x))^2 dx \right) \leq C n^{-2s(1-\delta)/(2s+\gamma+1)}.$$

The proof of Theorem 4.1 is complete. \square

Proof of Proposition 5.1. Using a standard covariance equality and (12), for any $g \in \mathbb{L}^1([0, 1])$ and any $m \in \{1, \dots, n\}$, we have

$$\begin{aligned}
& |\mathbb{C}_{ov}(g(S_m), g(S_0))| \\
&= \left| \int_0^1 \int_0^1 (f_{(S_m, S_0)}(x, y) - f_S(x)f_S(y))g(x)g(y)dx dy \right| \\
&\leq \int_0^1 \int_0^1 |f_{(S_m, S_0)}(x, y) - f_S(x)f_S(y)||g(x)||g(y)|dx dy \\
&\leq 2C_* \int_0^1 \int_0^1 |g(x)||g(y)|dx dy = 2C_* \left(\int_0^1 |g(x)|dx \right)^2. \quad (24)
\end{aligned}$$

Moreover, using $(\phi_{j,k})^{(u)}(x) = 2^{(2u+1)j/2}\phi^{(u)}(2^j x - k)$ and doing the change of variables $y = 2^j x - k$, we have

$$\begin{aligned}
& \int_0^1 |T_\nu(\phi_{j,k})(x)|dx \leq \nu! \sum_{u=0}^\nu \int_0^1 |x^u(\phi_{j,k})^{(u)}(x)|dx \\
&\leq \nu! \sum_{u=0}^\nu \int_0^1 |(\phi_{j,k})^{(u)}(x)|dx = C2^{-j/2} \sum_{u=0}^\nu 2^{uj} \int |\phi^{(u)}(y)|dy \\
&\leq C2^{\nu j} 2^{-j/2}. \quad (25)
\end{aligned}$$

It follows from (24) and (25) that

$$|\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_m), T_\nu(\phi_{j,k})(S_0))| \leq C2^{2\nu j} 2^{-j}. \quad (26)$$

By [8, equation before eq. (20)], we have

$$|\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C2^{(2\nu+q)j} \alpha_m^q.$$

Therefore

$$|\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C \min \left(2^{2\nu j} 2^{-j}, 2^{(2\nu+q)j} \alpha_m^q \right).$$

Hence, by (11),

$$\begin{aligned}
& \sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \\
&\leq C \sum_{m=1}^n \min \left(2^{2\nu j} 2^{-j}, 2^{(2\nu+q)j} \alpha_m^q \right) \\
&\leq C \left(\sum_{m=1}^{2^j-1} 2^{2\nu j} 2^{-j} + \sum_{m=2^j}^n 2^{(2\nu+q)j} \alpha_m^q \right) \\
&\leq C2^{2\nu j} \left(1 + \sum_{m=1}^n m^q \alpha_m^q \right) \leq C2^{2\nu j} (1 + n^\delta) \leq C2^{2\nu j} n^\delta.
\end{aligned}$$

Therefore

$$\sum_{k=0}^{2^j-1} \sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C2^{(2\nu+1)j}n^\delta.$$

Proposition 5.1 follows from Theorem 4.1 with $\gamma = 2\nu$. \square

Proof of Proposition 5.2. Since $(S_i)_{i \in \mathbb{Z}}$ is β -mixing, for any bounded function g , [32, equation line 12 p. 479 and Lemma 4.2. with $p = 1$] imply that

$$\sum_{m=0}^n |\mathbb{C}_{ov}(g(S_m), g(S_0))| \leq 2 \int_0^1 b(x)g^2(x)f_S(x)dx, \quad (27)$$

where b is a function such that, by (13), $\int_0^1 b(x)f_S(x)dx \leq \sum_{m=0}^\infty \beta_m \leq C$.

We have $(\phi_{j,k})^{(u)}(x) = 2^{(2u+1)j/2}\phi^{(u)}(2^jx - k)$. Since ϕ is compactly supported, we have $\sup_{x \in [0,1]} \sum_{k=0}^{2^j-1} (\phi^{(u)}(2^jx - k))^2 \leq C$. Therefore, for any integer $j \geq \tau$,

$$\begin{aligned} \sum_{k=0}^{2^j-1} (T_\nu(\phi_{j,k})(x))^2 &\leq \nu(\nu!)^2 \sum_{k=0}^{2^j-1} \sum_{u=0}^{\nu} x^{2u} ((\phi_{j,k})^{(u)}(x))^2 \\ &\leq \nu(\nu!)^2 \sum_{k=0}^{2^j-1} \sum_{u=0}^{\nu} ((\phi_{j,k})^{(u)}(x))^2 \\ &\leq \nu(\nu!)^2 2^{(2\nu+1)j} \sum_{u=0}^{\nu} \sum_{k=0}^{2^j-1} (\phi^{(u)}(2^jx - k))^2 \\ &\leq C2^{(2\nu+1)j}. \end{aligned} \quad (28)$$

Putting (27) and (28) together, we obtain

$$\begin{aligned} &\sum_{k=0}^{2^j-1} \sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \\ &\leq 2 \int_0^1 b(x)f_S(x) \sum_{k=0}^{2^j-1} (T_\nu(\phi_{j,k}))^2(x)dx \leq C2^{(2\nu+1)j} \int_0^1 b(x)f_S(x)dx \\ &\leq C2^{(2\nu+1)j}. \end{aligned}$$

Proposition 5.2 follows from Theorem 4.1 with $\gamma = 2\nu$. \square

Proof of Proposition 5.3. Since S_1, \dots, S_n are PPQD, the Newman inequality (see [23]) yields: for any $m \in \{1, \dots, n\}$ and any $g \in \mathcal{C}^1([0, 1])$,

$$|\mathbb{C}_{ov}(g(S_m), g(S_0))| \leq \left(\sup_{x \in [0, 1]} |g'(x)| \right)^2 \mathbb{C}_{ov}(S_m, S_0). \quad (29)$$

Moreover, since $(\phi_{j,k})^{(u)}(x) = 2^{(2u+1)j/2} \phi^{(u)}(2^j x - k)$, we have

$$\begin{aligned} \sup_{x \in [0, 1]} |T'_\nu(\phi_{j,k})(x)| &\leq (\nu + 1)! \sum_{u=0}^{\nu+1} \sup_{x \in [0, 1]} |x^u (\phi_{j,k})^{(u)}(x)| \\ &\leq C \sum_{u=0}^{\nu+1} 2^{(2u+1)j/2} \sup_{x \in [1-N, N]} |\phi^{(u)}(x)| \\ &\leq C 2^{(2(\nu+1)+1)j/2} = C 2^{(2\nu+3)j/2}. \end{aligned} \quad (30)$$

Therefore, by (26) (which holds thanks to (12)), (29) and (30), we have

$$|\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_m), T_\nu(\phi_{j,k})(S_0))| \leq C \min(2^{2\nu j} 2^{-j}, 2^{(2\nu+3)j} \mathbb{C}_{ov}(S_m, S_0)).$$

Hence, by (14),

$$\begin{aligned} &\sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \\ &\leq C \sum_{m=1}^n \min\left(2^{2\nu j} 2^{-j}, 2^{(2\nu+3)j} \mathbb{C}_{ov}(S_m, S_0)\right) \\ &\leq C \left(\sum_{m=1}^{2^j-1} 2^{2\nu j} 2^{-j} + \sum_{m=2^j}^n 2^{(2\nu+3)j} \mathbb{C}_{ov}(S_m, S_0) \right) \\ &\leq C 2^{2\nu j} \left(1 + \sum_{m=1}^n m^3 \mathbb{C}_{ov}(S_m, S_0) \right) \leq C 2^{2\nu j} (1 + n^\delta) \leq C 2^{2\nu j} n^\delta. \end{aligned}$$

Therefore

$$\sum_{k=0}^{2^j-1} \sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C 2^{(2\nu+1)j} n^\delta.$$

Proposition 5.3 follows from Theorem 4.1 with $\gamma = 2\nu$. \square

Proof of Proposition 5.4. Proceeding similarly to the proof of Proposition 5.3, we have

$$|\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_m), T_\nu(\phi_{j,k})(S_0))| \leq C 2^{(2\nu+3)j} \mathbb{C}_{ov}(S_m, S_0).$$

Therefore, by (16),

$$\sum_{k=0}^{2^j-1} \sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C2^{(2\nu+4)j} n^\delta.$$

Proposition 5.4 follows from Theorem 4.1 with $\gamma = 2\nu + 3$. \square

Proof of Proposition 5.5. A standard covariance inequality for ρ -mixing gives: for any $m \in \{1, \dots, n\}$ and any $g \in \mathbb{L}^2([0, 1], f_S(x)dx)$,

$$|\mathbb{C}_{ov}(g(S_m), g(S_0))| \leq \mathbb{E}((g(S_0))^2) \rho_m. \quad (31)$$

See, for instance, [35, Lemma 1.2.7.].

Since $S_0(\Omega) = [0, 1]$, for any integer $j \geq \tau$ and any $k \in \{0, \dots, 2^j - 1\}$, we have

$$\begin{aligned} \mathbb{E}((T_\nu(\phi_{j,k})(S_0))^2) &\leq \nu(\nu!)^2 \sum_{u=0}^{\nu} \mathbb{E}\left(S_0^{2u}((\phi_{j,k})^{(u)}(S_0))^2\right) \\ &\leq \nu(\nu!)^2 \sum_{u=0}^{\nu} \mathbb{E}\left(((\phi_{j,k})^{(u)}(S_0))^2\right). \end{aligned} \quad (32)$$

Using (17), $(\phi_{j,k})^{(u)}(x) = 2^{(2u+1)j/2} \phi^{(u)}(2^j x - k)$ and doing the change of variables $y = 2^j x - k$, we obtain

$$\begin{aligned} \mathbb{E}\left(((\phi_{j,k})^{(u)}(S_0))^2\right) &= \int_0^1 ((\phi_{j,k})^{(u)}(x))^2 f_S(x) dx \\ &\leq C_* \int_0^1 ((\phi_{j,k})^{(u)}(x))^2 dx \\ &\leq C_* 2^{2uj} \int (\phi^{(u)}(y))^2 dy. \end{aligned} \quad (33)$$

Putting (32) and (33) together, we obtain

$$\mathbb{E}((T_\nu(\phi_{j,k})(S_0))^2) \leq C_* \nu(\nu!)^2 \sum_{u=0}^{\nu} 2^{2uj} \int_{1-N}^N (\phi^{(u)}(y))^2 dy \leq C2^{2\nu j}. \quad (34)$$

Hence, by (31), (34) and (16),

$$\sum_{m=1}^n |\mathbb{C}_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C2^{2\nu j} \sum_{m=1}^n \rho_m \leq C2^{2\nu j} n^\delta.$$

Therefore

$$\sum_{k=0}^{2^j-1} \sum_{m=1}^n |C_{ov}(T_\nu(\phi_{j,k})(S_0), T_\nu(\phi_{j,k})(S_m))| \leq C2^{(2\nu+1)j}n^\delta.$$

Proposition 5.5 follows from Theorem 4.1 with $\gamma = 2\nu$. \square

Acknowledgments

We thank the referee for insightful comments that helped us improve the paper significantly.

References

- [1] Andersen, K. and Hansen, M. (2001), Multiplicative censoring: density estimation by a series expansion approach. *Journal of Statistical Planning and Inference*, **98**, 137-155.
- [2] Antoniadis, A. (1997), Wavelets in statistics: a review (with discussion). *Journal of the Italian Statistical Society, Series B*, **6**, 97-144.
- [3] Bradley, R. C. (2005), Basic Properties of Strong Mixing Conditions. A Survey and Some Open Questions, *Probability Surveys*, **2**, 107-144.
- [4] Cai, Z. W. and Roussas, G. G. (1997), Efficient estimation of a distribution function under quadrant dependence. *Scandinavian Journal of Statistics*, **24**, 1-14.
- [5] Carrasco, M. and Chen, X. (2002), Mixing and moment properties of various GARCH and stochastic volatility models. *Econometric Theory*, **18**, 17-39.
- [6] Chaubey, Y. P., Doosti, H., and Prakasa Rao, B. L. S. (2006), Wavelet based estimation of the derivatives of a density with associated variables. *International Journal of Pure and Applied Mathematics*, **27**(1), 97-106.
- [7] Chaubey, Y. P., Chesneau, C., and Doosti, H. (2011), On linear wavelet density estimation: some recent developments. *Journal of the Indian Society of Agricultural Statistics*, **65**(2), 169-179.

- [8] Chesneau, C. (2011), Wavelet estimation of a density in a GARCH-type model. *Communications in Statistics: Theory and Methods*, to appear.
- [9] Cohen, A., Daubechies, I., Jawerth, B., and Vial, P. (1993), Wavelets on the interval and fast wavelet transforms. *Applied and Computational Harmonic Analysis*, **24**(1), 54–81.
- [10] Comte, F., Dedecker, J., and Taupin, M.-L. (2008), Adaptive density estimation for general ARCH models. *Econometric Theory*, **24**(6), 1628-1662.
- [11] Dewan, I. and Prakasa Rao, B. L. S. (1999), A general method of density estimation for associated random variables. *Journal of Nonparametric Statistics*, **10**, 405-420.
- [12] Doukhan, P. (1994), *Mixing. Properties and Examples*. Lecture Notes in Statistics, **85**, New York: Springer Verlag.
- [13] Doukhan, P. and Louhichi, S. (1998), Estimation de la densité d'une suite faiblement dépendante. *C. R. Acad. Sci. Paris Sr. I Math.*, **327**(12), 989-992.
- [14] Doukhan, P. and Louhichi, S. (1999), A new weak dependence condition and applications to moment inequalities. *Stochastic Processes and their Applications*, **84**, 313-342.
- [15] Fryzlewicz, P. and Subba Rao, S. (2011), Mixing properties of ARCH and time-varying ARCH processes. *Bernoulli*, **17**(1), 320-346.
- [16] Härdle, W., Kerkycharian, G., Picard, D., and Tsybakov, A. (1998), *Wavelet, Approximation and Statistical Applications*. Lectures Notes in Statistics, **129**, New York: Springer Verlag.
- [17] Hosseinioun, N., Doosti, H., and Nirumand, H. A. (2011), Nonparametric estimation of the derivatives of a density by the method of wavelet for mixing sequences. *Statistical Papers*, to appear.
- [18] Kolmogorov, A. N. and Rozanov, Yu. A. (1960), On strong mixing conditions for stationary Gaussian processes. *Theor. Probab. Appl.*, **5**, 204-208.
- [19] Leblanc, F. (1996), Wavelet linear density estimator for a discrete time stochastic process: L_p -losses. *Statistics and Probability Letters*, **27**, 71-84.

- [20] Lehmann, E. L. (1966), Some concepts of dependence. *The Annals of Mathematical Statistics*, **37**, 1137-1153.
- [21] Masry, E. (2001), Multivariate probability density estimation for associated processes: Strong consistency and rates. *Statistics and Probability Letters*, **58**, 205-219.
- [22] Meyer, Y. (1992), *Wavelets and Operators*. Cambridge: Cambridge University Press.
- [23] Newman, C. M. (1980), Normal fluctuations and the FKG inequalities. *Commun. Math. Phys.*, **74**, 119-128.
- [24] Prakasa Rao, B. L. S. and Dewan, I. (2001), Associated sequences and related inference problems. *Handbook of Statistics*, **19**, Stochastic Processes: Theory and Methods, (eds. C.R. Rao and D.N. Shanbag), 693-728, North Holland, Amsterdam.
- [25] Prakasa Rao, B. L. S. (2003), Wavelet linear density estimation for associated sequences. *Journal of the Indian Statistical Association*, **41**, 369-379.
- [26] Rosenblatt, M. (1956), A central limit theorem and a strong mixing condition. *Proc. Nat. Ac. Sc. USA*, **42**, 43-47.
- [27] Roussas, G. G. (1999), Positive and negative dependence with some statistical applications. *Asymptotics, nonparametrics and time series* (ed. S. Ghosh), 757-788, Marcel Dekker, New York.
- [28] Sancetta, A. (2009), Strong law of large numbers for pairwise positive quadrant dependent random variables. *Statistical Inference for Stochastic Processes*, **12**(1), 55-64.
- [29] Shao, Q.-M. (1995), Maximal inequality for partial sums of ρ -mixing sequences. *Ann. Probab.*, **23**, 948-965.
- [30] Tribouley, K. and Viennet, G. (1998), L_p adaptive density estimation in a β -mixing framework. *Ann. Inst. H. Poincaré Probab. Statist.*, **34**, 179-208.
- [31] Van Zanten, H. and Zareba, P. (2008), A note on wavelet density deconvolution for weakly dependent data. *Stat. Inference Stoch. Process.*, **11**, 207-219.

- [32] Viennet, G. (1997), Inequalities for absolutely regular processes: application to density estimation. *Probab. Theory Related Fields*, **107**, 467-492.
- [33] Volkonskii, V. A. and Rozanov, Y. A. (1959), Some limit theorems for random functions I. *Theory Probab. Appl.*, **4**, 178-197.
- [34] Withers, C. S. (1981), Conditions for linear processes to be strong-mixing. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, **57**, 477-480.
- [35] Zhengyan, L. and Lu, C. (1996), *Limit Theory for Mixing Dependent Random Variables*. Kluwer, Dordrecht.

